



Is Alvarado Score Helpful for Pediatrician in Diagnosis of Acute Appendicitis? Our Center Experience

Sedigheh Rafiei Tabatabaei ¹, Abdollah Karimi ¹, Mohammad Nassiri ¹, Leily Mohajerzadeh ^{1,2,*}, Shahnaz Armin ¹, Roxana Mansour Ghanaie ¹, Seyed Alireza Fahimzad ¹, Farideh Shiva ¹, Fatemeh Fallah ¹, Ahmad Reza Shamshiri ¹, Iraj Sedighi ³, Shirin Sayyahfar ⁴, Mohammad Sadegh Rezai ⁵, Babak Abdinia ⁶, Hosein Heydari ⁷, Gholamreza Soleimani ⁸, Mostafa Behpour Oskouee ⁹, Jafar Soltani ¹⁰, Mehran Ahmadi ¹¹, Manijeh Kabhazi ¹² and Sayeh Hatefi ²

¹Pediatric Infections Research Center (PIRC), Research Institute for Children's Health (RICH), Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Pediatric Surgery Research Center, Research Institute for Children's Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Pediatric Department, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

⁴Research Center of Pediatric Infectious Disease, Institute of Immunology and Infectious Disease, Iran University of Medical Sciences, Tehran, Iran

⁵Pediatric Infectious Disease Research Center, Communicable Disease Institute, Mazandaran University of Medical Sciences, Sari, Iran

⁶Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

⁷Research Center, Qom University of Medical Sciences, Qom, Iran

⁸Children and Adolescent Health Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

⁹Department of Pediatrics, Najmieh Hospital, Baghyatollah University of Medical Sciences, Tehran, Iran

¹⁰Department of Pediatrics, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

¹¹Clinical Research Development Center of Children Hospital, Hormozgan University of Medical Sciences, Hormozgan, Iran

¹²Infectious Diseases Research Center (IDRC), Arak University of Medical Sciences, Arak, Iran

*Corresponding author: Pediatric Surgery Research Center, Research Institute for Children's Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: mohajerzadehl@yahoo.com

Received 2020 August 24; Revised 2020 December 07; Accepted 2020 December 11.

Abstract

Background: Many problems in the diagnosis of patients with suspected appendicitis have led to the design of clinical scoring systems. In children, diagnostics tools for appendicitis are more critical. Younger patients, diagnostic challenges become more. Practical scoring systems are useful without any particular material and necessitate novel ability.

Objectives: However, in spite of the reported outstanding consequences, these scoring systems are not employed regularly.

Methods: In this cross-sectional study performed from October 2016 to October 2017, ten provinces out of the 31 provinces in Iran were randomly selected. A total of 631 patients referring to the hospitals with the suspicion of acute appendicitis were assessed. Related variables such as age, sex, right lower quadrant (RLQ) pain, migration of pain to RLQ, nausea, and presence of vomiting, anorexia, tenderness in RLQ and guarding, presence of rebound tenderness, and degree of fever were taken from recorded files by pediatricians. Alvarado scoring system was used for included cases to assess the accuracy of this test for diagnosis of appendicitis in our centers.

Results: The mean age of eligible patients was 9.3 ± 3.21 years ranged from 3 years to 18 years, and 380 (60.5%) were male. Using the Alvarado score system in this study, considering the cut-off point value of 7 to decide for operation, the positive predictive value (PPV) showed 32.6%, and negative predictive value (NPV) was 76.73%, with a sensitivity of 44.05% and, specificity of 66.95%. There were statistically no significant correlations among the scoring of the Alvarado and diagnosis of AA ($P < 0.05$). Regarding receiver operator characteristic curves (ROC), the area under curve (AUC) was 0.58 (0.54 to 0.63) for Alvarado. The AUC was very low, so there was no value for the diagnosis of appendicitis. According to the findings of the present study, the cut-off point of 4.5 is suggested for the diagnosis of acute appendicitis in children with a sensitivity of 73% and specificity of 58%.

Conclusions: Although the Alvarado scores supply obviously practical diagnostic information in the management of pediatric population with supposed appendicitis, this method delivers no adequate PPV for clinical practice as a safe way for determining necessary operation.

Keywords: Acute Appendicitis, Alvarado Score, Abdominal Tenderness

1. Background

Acute appendicitis (AA) is the most frequent cause of emergency conditions of the abdominal cavity. However, unfortunately, appendicitis is still an intricate illness en-

tity (1). Regarding the lack of widespread investigations in this entity, there are several vague points for clinicians worldwide (2). Diagnostic errors for appendicitis are relatively common in childhood, thereby resulting in a delay for surgery or negative appendectomies.

Acute appendicitis is uncommon in children younger than 3 years, but the rate of perforation is much higher in young children.

Regarding the importance of AA diagnosis in children and the presence of numerous misdiagnoses for this condition, there are some scoring systems for early identification of acute abdomen in the literature. We used the Alvarado scoring system (3) to evaluate the value of this diagnostic way in improving preoperative diagnosis of AA. Alvarado scoring system identifies pediatric patients with suspected appendicitis. Parameters for evaluation of appendicitis in the Alvarado score include migration of pain, anorexia, nausea or vomiting, right lower abdominal quadrant tenderness, rebound tenderness in the right lower quadrant, increase of temperature, leukocytosis, shift to the left of neutrophils.

2. Objectives

We aimed to investigate retrospectively the diagnostic effectiveness of the preoperative tests based on histopathologic consequences.

3. Methods

In this cross-sectional study performed from October 2016 to October 2017, ten provinces out of 31 provinces in Iran were randomly selected. A total of 631 patients referring to the specified hospitals with the suspicion of acute appendicitis were assessed. The data were gathered and entered into the checklist by reviewing the hospital recorded files.

3.1. Selection of Patients for Study and Data Collection

This study was designed for Children between 3 to 18 years of age, with acute abdominal pain suspected of appendicitis that at first were visited by a pediatrician in the pediatric emergency medical center and then required surgical consultation for possible appendicitis before request radiologic investigations. Cases with a history of previous abdominal surgery who had chronic medical conditions, or underwent multiple radiologic investigations of the abdomen within the previous last week were excluded. About 20 cases were excluded. A total of 631 patients who were referred to the specified hospitals with the suspicion of acute appendicitis and had surgical consultation for probable appendicitis were assessed. The data were gathered and entered into the checklist by reviewing the hospital recorded files.

The patient's medical records were investigated for symptoms and signs and physical examination recorded by a pediatrician and all data registered based on a non-surgeon examiner who consulted with a surgeon. Laboratory, pathology, and operative reports were noted frankly from the electronic laboratory information. Operative information was evaluated for the occurrence or lack of appendiceal perforation. Reports of pathology were analyzed whether there was appendicitis. Appendix pathology reports such as "early appendicitis", "appendicitis", "gangrenous", "perforated appendix", all of which were considered appendicitis. Another impression registered was the absence of appendicitis. Cases with missing data were excluded. The manuscript was accepted by the Institutional Board Review and the Medical Ethics Committee of Shahid Beheshti University of Medical Sciences.

3.2. Methods of Outcome Measurement

The final diagnosis was confirmed either by pathologic details if the patient underwent an operation in the same center or by follow-up telephone calls if the patient did not have an operation history. Telephone call revealed further diagnostic evaluation or operative management in another center for two weeks after the initial visit to the primary center. Data used for the assessment of Alvarado's scoring system (3) are summarized in Table 1. Alvarado scoring system was used to include cases for the assessment of the accuracy of this test for diagnosing appendicitis in our centers.

Table 1. Alvarado Scoring System for Pediatric Appendicitis (3)

Characteristic	Point Assessment	Suggested Management
1. Migration of pain	1	
2. Anorexia	1	
3. Nausea/vomiting	1	
4. RLQ tenderness	1	
5. Rebound pain	2	
6. Fever (> 37.3)	1	
7. Leukocytosis (L) (10,000/ μ L)	2	
8. Left shift: neutrophil > 75%	1	
Points of Alvarado Scoring system		
5 - 6 likelihood of appendicitis		Observe
Sum of 7 and more suggests appendicitis		Surgery

3.3. Data Analysis

Data were analyzed with SPSS version 17. Patients with missing data (for any component of the Alvarado scoring system) were removed. The computed scores were employed to evaluate the Alvarado score greater than or equal to seven for establishing risk for appendicitis. In addition, receiver operator characteristic curves (ROC) were used for all scores to graphically characterize ROC curves and evaluation of the diagnostic value of the documented parameters in acute appendicitis.

Data were measured with student *t*-test and chi-square test for acute appendicitis and non-appendicitis groups. Quantitative data were described as mean and standard deviation (SD), which were compared with *t*-test. Qualitative data were compared with the chi-square test with frequency and percentage of description. A P-value of less than 0.05 was considered statistically significant.

4. Results

4.1. Demographic Characteristics of the Patients

The mean age of eligible patients was 9.3 ± 3.21 years ranged from three to 18 years, of which 380 (60.5%) were male. During the period of October 2014 and October 2015, 631 patients had a surgical consultation for probable appendicitis among patients with abdominal pain. Of all 631 patients who were suspected of appendicitis, definitive diagnosis was confirmed in 168 with the prevalence rate of 27% that perforated appendicitis was found in 22 patients, infectious appendicitis in 14 patients, and gangrenous appendicitis in 11 patients. The mean age of patients without appendicitis was 108 ± 38.81 months, and the mean age of patients with acute appendicitis (AA) was 111 ± 39.92 months.

4.2. Patients' Symptoms and Signs

Migratory pain was present in 41% of patients without appendicitis and in 53.2% of patients with AA (P-value = 0.040). In this study, there were no significant differences for migratory pain between the two groups. Nausea or vomiting was noticed in 70% of patients without appendicitis and 75% of patients with AA (P-value = 0.392). In this study, there were no significant differences in nausea or vomiting between the two groups. Anorexia existed in 62% of patients without appendicitis and was present in 70% of children with AA (P-value = 0.069). In this study, there were no significant differences for anorexia between the two groups.

4.3. Patient's Physical Examination

Tenderness in the right iliac fossa was found in 42% of the patients without appendicitis and was found in 67% of the patients with AA (P-value = 0.000). In this study, there were no significant differences for tenderness in RLQ between the two groups. Rebound tenderness in the right iliac fossa was found in 42% of patients without appendicitis, and it was found in 68% of the patients with AA (P-value = 0.000). In this study, there were no significant differences for Rebound tenderness between the two groups. An increase in auxiliary temperature $> 37.3^{\circ}\text{C}$ was noticed in 2.35% of the patients without appendicitis and was present in 3.5% of the children with AA (P-value = 0.386). In this study, there were no significant differences for fever between the two groups.

4.4. Laboratory Findings

Leukocytosis $> \text{WBC} = 10,000/\text{L}$ was noticed in 71% of patients without appendicitis and was present in 72.8% of children with AA (P-value = 0.629). In this study, there were no significant differences for leukocytosis between the two groups. Left shift with neutrophil $> 75\%$ was presented in 60.2% of the patients without appendicitis and was present in 50% of the children with AA (P-value = 0.016). Moreover, there were no significant differences for shift to the left between the two groups. Clinical characteristics of the cases with and without appendicitis are summarized in [Table 2](#).

4.5. Alvarado Scores

Alvarado scores for Appendicitis group: anorexia (70% of patients), nausea/vomiting (75%), RLQ tenderness (67.3%), differential WBC count (50%), migration of pain (53.2%), rebound pain (66.5%), presence of fever (3.5%), and increase in leukocyte count (72.8%). When the Alvarado system was used, regarding the cut-off point of 7 to decide for operation, the positive predictive value (PPV) was 32.6% and negative predictive value (NPV) was 76.73%, with sensitivity of 44.05% and, specificity of 66.95%. See [Table 3](#) for sensitivity, specificity, NPV, PPV, accuracy, and the ratio for clinical Alvarado scoring systems.

In this study, there were no significant differences between the values of the Alvarado score and the occurrence of AA. Also, ROC curves were created for Alvarado scores to symbolize the whole score presentation ([Figure 1](#)). The area under the curve was 0.58 (0.54 to 0.63) for Alvarado. The area under the curve is very low; so there is no value for diagnosis of appendicitis.


All probable cut-off points, with their related sensitivity and false-positive rate, are shown in [Table 4](#). Scores of

Table 2. Clinical Characteristics of the Patients with Suspected Appendicitis

Feature (Total = 631)	No Appendicitis, N = 463 (73)	Appendicitis, N = 168 (27)	P-Value
Mean age, mo	108 ± 38.81	111 ± 39.92	0.000
Migration of pain, %	41	53.2	0.040
Anorexia, %	62	70	0.069
Nausea/vomiting, %	70	75	0.392
RLQ tenderness, %	42	67	0.000
Rebound pain, %	46	66.5	0.000
Increase in temperature > 37.3°C, %	2.3	3.5	0.386
Leukocytosis > WBC = 10,000/ μ L, %	71	72.8	0.629
Shift to the left, %	60.2	50	0.016

Table 3. Presentation of the Alvarado Scores More Than or Equal to 7

Alvarado	All Patients (N = 661)
Sensitivity, %	44.05
Specificity, %	66.95
Accuracy, %	60.86
Negative predictive value (NPV), %	76.73
Positive predictive value (PPV), %	32.60

**Figure 1.** Receiver operator characteristic curves (ROC) for arrangement of the Alvarado Scoring systems.

7 (Alvarado) have been determined in studies to discriminate cut-off points for diagnosing appendicitis. In the present study, the cut-off point of 4.5 was considered for children with a sensitivity of 73% and specificity of 58%.

Other diagnosis have been included: Mesenteries lym-

Table 4. Demonstrates Cut-off Points for All Probable Scores

Positive if Greater Than or Equal To	Sensitivity	1-Specificity
1.00	1.000	1.000
0.50	1.000	0.970
1.50	0.976	0.916
2.50	0.929	0.840
3.50	0.839	0.704
4.50	0.726	0.579
5.50	0.583	0.449
6.50	0.440	0.330
7.50	0.327	0.251
8.50	0.119	0.134
9.50	0.006	0.002
11.00	0.000	0.000

phadenitis, Meckel's diverticulitis, abscess of the ovary, torsion of ovarian cyst, ruptured ovarian cyst, and two cases without any recognized anomaly.

5. Discussion

Appendicitis is a prevalent disorder with a chance of occurrence about 7 percent for everybody in all lifetime (4) the most important symptoms are pain in the right lower quadrant, anorexia, and vomiting. The reliable physical exams are tenderness, rebound tenderness, and guarding in the iliac fossa. Practical investigations for suspicious appendicitis include complete blood count and urinalysis. Although the main diagnostic way remains physical examination, the use of imaging studies is growing more than before because of frequent misdiagnosis, especially in young patients (5). Recent studies showed that in-

flammatory markers, including white blood cell (WBC), C-reactive protein (CRP), or pro-calcitonin are valuable to recognize appendicitis with significant sensitivity and specificity (6, 7).

Unfortunately, the vague signs and symptoms in young children lead to increasing the risk of appendice perforation and related morbidity and mortality (8). Perforation of appendix reported 20% to 90% in various studies (9). The rate of negative appendectomy reported about 20%. So, to diagnose appendicitis, especially in the pediatric group, imaging can be a critical tool for decreasing misdiagnosis (9, 10). Ultrasonography has been verified as an early diagnostic tool in young patients to avoid radiation exposure and risk of cancer secondary to CT (11, 12). Routinely, CT scan results in extensive cost too.

Scoring systems are simple and can be used independently of particular tools, and they do not require novel abilities (7). On the other hand, these scorings are not employed regularly, so the results are not clear, especially in children. In different scoring systems for judgment of AA, the Alvarado is shown to be more successful. A score of more than seven points is reported with a sensitivity of about 88% to 90%, but the specificity is usually under 80% and is particularly low in young patients and women (7). Alvarado at first manuscript (3) retrospectively studied 305 cases with acute abdominal pain indicative of appendicitis. The tests showed sensitivity of 81%, specificity of 74%, PPV of 92%, and NPV of 46%. In the present study, 631 patients, aged 3 to 18 years, with suspected appendicitis were assessed. Twenty-seven percent of cases had appendicitis. Moreover, PPV was 32.6%, and NPV was 76.73%, with a sensitivity of 44.05% and specificity of 66.95%.

Ohle et al. (13) performed a review article and evaluated the diagnostic accuracy of the Alvarado score. The investigation was designed with paying attention to three sub-groups: women, men, and children. They suggested the cut-off point of five is suitable for 'ruling out' appendicitis (sensitivity 99% overall, 96% in men cases, 99% for woman cases, and 99% in children). The cut-off point of seven, as suggested by Alvarado for the need to attempt surgery, had poor results for each subgroup (with specificity overall about 81%, for men 57%, women cases 73%, and children cases 76%). Similarly, in the present study, 4.5 is suggested as the cut-off point for children with a sensitivity of 73% and specificity of 58%.

So it is better that a cut-off point score is used for predicting the likelihood of appendicitis in high-risk groups such as children lower than seven. As studies showed a high chance of perforation followed appendicitis in chil-

dren, so we suggest that the Alvarado score is used at the cut-off point of 4.5 to 5 to establish a rule-out point for appendicitis in children for preventing from misdiagnosis and catastrophic morbidity and mortality in children. However, we recommend the use of other investigation tools for proper diagnosis to prevent negative operation in young children.

Singla et al. (14) designed a cohort study on the Asian population for two years. They studied 50 adult cases with Right lower quadrant pain and used a modified Alvarado score (modified Alvarado score excludes shift to the left for neutrophil maturation more than 75%). Also, PPV was reported 100 %, and the rate of negative appendectomy was 0% (no false-positive patients, while 21 false-negative patients). In contrast, Khan and ur Rehman study (15) demonstrated a PPV of 83.3% with 15.6% negative appendectomy.

Goyal et al. (16) designed a study on patients supposed to acute appendicitis through one year by Modified Alvarado score (as told above). The study explained making use of the Modified Alvarado Score in cases with suspicion of appendicitis with an acceptable amount of sensitivity and specificity. They advised Modified Alvarado scores in men and children; however, supplementary employing of ultrasonography or diagnostic laparoscopy in women is suggested to diminish the incorrect rate of false-positive in them. However, in our study, shift to the left with neutrophil > 75% was presented in 60.2% of the patients without appendicitis and was present in 50% of the children with AA (P-value = 0.016). We have a statistically significant difference in the occurrence of shift to the left in the patients with AA compared with the cases without AA. Therefore, we do not recommend a modified Alvarado Score in children.

Toprak et al. (17) assessed the combination of Alvarado score with ultrasonography results in children suspected of appendicitis. They achieved 92% specificity, 93% sensitivity, and 92% accuracy for the US in diagnosis of appendicitis; but confirmed a few obstacles and limits for the US that led to false-positive or -negative consequences. So they employed Alvarado score integration with the US according to recognize the chance of appendicitis in children. They achieved a cut-off of 6 for Alvarado score as a good predictor of appendicitis in children. All patients with an Alvarado score \geq of 7 had a diagnosis of acute appendicitis. The present study proposes that if the Alvarado score is more than 6 scores and US does not show any compressible appendix, surgery is suitable management without the need for additional imaging techniques, particularly CT. They recommended that in the patients with an ab-

sence of none compressible appendix with a low Alvarado score, appendicitis can be securely ruled out. Further, CT is suggested to be helpful in children with moderate scores and vague US results (18-20). The negative appendectomy and complicated appendicitis rates are both vital for superiority measures of the management of appendicitis, especially in children (21, 22). On the other hand, these two items have an opposite correlation. In our study, as the cut-off point of 7 for the Alvarado score is not suitable for suggestion of appendicitis, appropriate diagnosis needs other investigations accompanied with Alvarado score.

5.1. Conclusions

In summary, although impression to clinical Alvarado score system can be an influential guideline in decision for cases with acute abdomen suspected of appendicitis in adults, judgment on examination of abdomen is complex in children, especially in young children. Furthermore, although these scores are attractive in conception, pediatric clinicians- in the fashion of deciding on the diagnosis of the disease in children- should observe the precautionary measures and examine them very comprehensively. Furthermore, although the Alvarado system scoring supplies obviously helpful diagnostic information in adult patients but in children supposed to appendicitis, this method lonely affords not sufficient positive predictive value (PPV) for determining the necessitate of operation. So it is healthier that cut off point score for expecting the possibility of appendicitis in children judge lower than 7.

Footnotes

Authors' Contribution: Study concept and design: Sedigheh Rafiei Tabatabaei, Abdollah Karimi, and Leily Mohajerzadeh. Acquisition of data: Shahnaz Armin, Roxana Mansour Ghanaie, Seyed Alireza Fahimzad, Farideh Shiva, Fatemeh Fallah, Iraj Sedighi, Shirin Sayyahfar, Mohammad Sadegh Rezai, Babak Abdinia, Hosein Heydar, Gholamreza Soleimani, Mostafa Behpour Oskouee, Jafar Soltani, Mehran Ahmadi, and Manijeh Kahbazi. Analysis and interpretation of data and statistical analysis: Ahmad Reza Shamshiri. Drafting of the manuscript: Leily Mohajerzadeh.

Conflict of Interests: The authors have no conflicts of interest to disclose.

Ethical Approval: This study was approved by in 8th session of the Organizational Committee of Ethics in Biomed-

ical Research of Shahid Beheshti University of Medical Sciences on 1394.07.12.

Funding/Support: This study is supported by Pediatric Infections Research Center (PIRC), Research Institute for Children's Health (RICH), Shahid Beheshti University of Medical Sciences, Tehran, Iran.

References

1. Richardsen I, Schob DS, Ulmer TF, Steinau G, Neumann UP, Klink CD, et al. Etiology of Appendicitis in Children: The Role of Bacterial and Viral Pathogens. *J Invest Surg.* 2016;**29**(2):74-9. doi: [10.3109/08941939.2015.1065300](https://doi.org/10.3109/08941939.2015.1065300). [PubMed: [26376211](https://pubmed.ncbi.nlm.nih.gov/26376211/)].
2. Alegbeleye Bamidele J. Current trends of acute appendicitis in africa: A clinical review. *J Int Healthcare Sci.* 2019;**7**(2):63-92.
3. Alvarado A. A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med.* 1986;**15**(5):557-64. doi: [10.1016/s0196-0644\(86\)80993-3](https://doi.org/10.1016/s0196-0644(86)80993-3). [PubMed: [3963537](https://pubmed.ncbi.nlm.nih.gov/3963537/)].
4. Subotic AM, Sijacki AD, Dugalic VD, Antic AA, Vukovic GM, Vukojevic VS, et al. Evaluation of the Alvarado score in the diagnosis of acute appendicitis. *Acta Chir Jugosl.* 2008;**55**(1):55-61. doi: [10.2298/aci0801055s](https://doi.org/10.2298/aci0801055s). [PubMed: [18510062](https://pubmed.ncbi.nlm.nih.gov/18510062/)].
5. Memisoglu K, Karip B, Mestan M, Onur E. The value of preoperative diagnostic tests in acute appendicitis, retrospective analysis of 196 patients. *World J Emerg Surg.* 2010;**5**:5. doi: [10.1186/1749-7922-5-5](https://doi.org/10.1186/1749-7922-5-5). [PubMed: [20181221](https://pubmed.ncbi.nlm.nih.gov/20181221/)]. [PubMed Central: [PMC2834661](https://pubmed.ncbi.nlm.nih.gov/PMC2834661/)].
6. Yu CW, Juan LI, Wu MH, Shen CJ, Wu JY, Lee CC. Systematic review and meta-analysis of the diagnostic accuracy of procalcitonin, C-reactive protein and white blood cell count for suspected acute appendicitis. *Br J Surg.* 2013;**100**(3):322-9. doi: [10.1002/bjs.9008](https://doi.org/10.1002/bjs.9008). [PubMed: [23203918](https://pubmed.ncbi.nlm.nih.gov/23203918/)].
7. Leeuwenburgh MM, Stockmann HB, Bouma WH, Houdijk AP, Verhagen MF, Vrouenraets B, et al. A simple clinical decision rule to rule out appendicitis in patients with nondiagnostic ultrasound results. *Acad Emerg Med.* 2014;**21**(5):488-96. doi: [10.1111/acem.12374](https://doi.org/10.1111/acem.12374). [PubMed: [24842498](https://pubmed.ncbi.nlm.nih.gov/24842498/)].
8. Farzal Z, Farzal Z, Khan N, Fischer A. The diagnostic dilemma of identifying perforated appendicitis. *J Surg Res.* 2015;**199**(1):164-8. doi: [10.1016/j.jss.2015.04.058](https://doi.org/10.1016/j.jss.2015.04.058). [PubMed: [25976856](https://pubmed.ncbi.nlm.nih.gov/25976856/)].
9. Srinivasan A, Servaes S, Pena A, Darge K. Utility of CT after sonography for suspected appendicitis in children: integration of a clinical scoring system with a staged imaging protocol. *Emerg Radiol.* 2015;**22**(1):31-42. doi: [10.1007/s10140-014-1241-1](https://doi.org/10.1007/s10140-014-1241-1). [PubMed: [24917390](https://pubmed.ncbi.nlm.nih.gov/24917390/)].
10. Soldo I, Radisic Biljak V, Bakula B, Bakula M, Simundic AM. The diagnostic accuracy of clinical and laboratory parameters in the diagnosis of acute appendicitis in the adult emergency department population - a case control pilot study. *Biochem Med (Zagreb).* 2018;**28**(3):30712. doi: [10.11613/BM.2018.030712](https://doi.org/10.11613/BM.2018.030712). [PubMed: [30429680](https://pubmed.ncbi.nlm.nih.gov/30429680/)]. [PubMed Central: [PMC6214693](https://pubmed.ncbi.nlm.nih.gov/PMC6214693/)].
11. Miglioretti DL, Johnson E, Williams A, Greenlee RT, Weinmann S, Solberg LI, et al. The use of computed tomography in pediatrics and the associated radiation exposure and estimated cancer risk. *JAMA Pediatr.* 2013;**167**(8):700-7. doi: [10.1001/jamapediatrics.2013.311](https://doi.org/10.1001/jamapediatrics.2013.311). [PubMed: [23754213](https://pubmed.ncbi.nlm.nih.gov/23754213/)]. [PubMed Central: [PMC3936795](https://pubmed.ncbi.nlm.nih.gov/PMC3936795/)].
12. Mathews JD, Forsythe AV, Brady Z, Butler MW, Goergen SK, Byrnes GB, et al. Cancer risk in 680 000 people exposed to computed tomography scans in childhood or adolescence: data linkage study of 11 million Australians. *BMJ.* 2013;**346**.
13. Ohle R, O'Reilly F, O'Brien KK, Fahey T, Dimitrov BD. The Alvarado score for predicting acute appendicitis: a systematic review. *BMC Med.*

- 2011;**9**:139. doi: [10.1186/1741-7015-9-139](https://doi.org/10.1186/1741-7015-9-139). [PubMed: [22204638](https://pubmed.ncbi.nlm.nih.gov/22204638/)]. [PubMed Central: [PMC3299622](https://pubmed.ncbi.nlm.nih.gov/PMC3299622/)].
14. Singla A, Singla S, Singh M, Singla D. A comparison between modified Alvarado score and RIPASA score in the diagnosis of acute appendicitis. *Updates Surg.* 2016;**68**(4):351-5. doi: [10.1007/s13304-016-0381-0](https://doi.org/10.1007/s13304-016-0381-0). [PubMed: [27338243](https://pubmed.ncbi.nlm.nih.gov/27338243/)].
 15. Khan I, ur Rehman A. Application of alvarado scoring system in diagnosis of acute appendicitis. *J Ayub Med Coll Abbottabad.* 2005;**17**(3):41-4. [PubMed: [16320795](https://pubmed.ncbi.nlm.nih.gov/16320795/)].
 16. Goyal DP, Kanwar DA, Malhotra DP, Sharma DD, Sharma DSK. Role of Modified Alvarado Score in the Diagnosis of Acute Appendicitis. *IOSR J Dent Med Sci.* 2014;**13**(10):44-8. doi: [10.9790/0853-131034448](https://doi.org/10.9790/0853-131034448).
 17. Toprak H, Kilincaslan H, Ahmad IC, Yildiz S, Bilgin M, Sharifov R, et al. Integration of ultrasound findings with Alvarado score in children with suspected appendicitis. *Pediatr Int.* 2014;**56**(1):95-9. doi: [10.1111/ped.12197](https://doi.org/10.1111/ped.12197). [PubMed: [23937630](https://pubmed.ncbi.nlm.nih.gov/23937630/)].
 18. Nasiri S, Mohebbi F, Sodagari N, Hedayat A. Diagnostic values of ultrasound and the Modified Alvarado Scoring System in acute appendicitis. *Int J Emerg Med.* 2012;**5**(1):26. doi: [10.1186/1865-1380-5-26](https://doi.org/10.1186/1865-1380-5-26). [PubMed: [22673121](https://pubmed.ncbi.nlm.nih.gov/22673121/)]. [PubMed Central: [PMC3410771](https://pubmed.ncbi.nlm.nih.gov/PMC3410771/)].
 19. Tan WJ, Acharyya S, Goh YC, Chan WH, Wong WK, Ooi LL, et al. Prospective comparison of the Alvarado score and CT scan in the evaluation of suspected appendicitis: a proposed algorithm to guide CT use. *J Am Coll Surg.* 2015;**220**(2):218-24. doi: [10.1016/j.jamcollsurg.2014.10.010](https://doi.org/10.1016/j.jamcollsurg.2014.10.010). [PubMed: [25488354](https://pubmed.ncbi.nlm.nih.gov/25488354/)].
 20. Wu HP, Chen CY, Kuo IT, Wu YK, Fu YC. Diagnostic values of a single serum biomarker at different time points compared with Alvarado score and imaging examinations in pediatric appendicitis. *J Surg Res.* 2012;**174**(2):272-7. doi: [10.1016/j.jss.2011.01.047](https://doi.org/10.1016/j.jss.2011.01.047). [PubMed: [21529827](https://pubmed.ncbi.nlm.nih.gov/21529827/)].
 21. Mohajerzadeh L, Rouzrokh M, Khaleghnejad Tabari A, Mirshemirani A, Atqiaee K, Dara N. Laparoscopic Appendectomy in Complicated Appendicitis of Children. *Ann Colorectal Res.* 2014;**2**(1). doi: [10.17795/acr-16599](https://doi.org/10.17795/acr-16599).
 22. Awayshih MMA, Nofal MN, Yousef AJ. Evaluation of Alvarado score in diagnosing acute appendicitis. *Pan Afr Med J.* 2019;**34**:15. doi: [10.11604/pamj.2019.34.15.17803](https://doi.org/10.11604/pamj.2019.34.15.17803). [PubMed: [31762884](https://pubmed.ncbi.nlm.nih.gov/31762884/)]. [PubMed Central: [PMC6859007](https://pubmed.ncbi.nlm.nih.gov/PMC6859007/)].